

LMU-TPW 95-8

June, 1995

Quark Mass Matrices in Orbifold Models

Tatsuo Kobayashi *

Sektion Physik, Universität München,
Theresienstr. 37, D-80333 München, Germany

Abstract

We study left-right symmetric quark mass matrices whose up- and down-sectors have the same structure. This type of realistic mass matrices are derived from orbifold models.

*Alexander von Humboldt Fellow

e-mail: kobayash@lswes8.ls-wess.physik.uni-muenchen.de

The origin of the fermion masses is one of the most important problems in particle physics. Higher dimension couplings could explain the hierarchical structure of the fermion masses and mixing angles [1, 2, 3, 4]. Underlying symmetries provide selection rules for higher dimension couplings as well as renormalizable couplings. In this case entries in mass matrices are written by suppression factors with powers, which are determined in terms of some types of quantum numbers under underlying symmetries. If we assume the left-right symmetry, the quark mass matrices are dominated by quantum numbers of the quark doublets Q_i as well as the Higgs fields. Hence from the assumption of the left-right symmetry it is plausible to expect that the quark mass matrix of the up-sector has the same structure as one of the down-sector. In refs.[3, 4], one type of up-down symmetric quark mass matrices are obtained by using an extra U(1) symmetry.

Superstring theory is the only known candidate for the unified theory of all the interactions including gravity. In superstring theory selection rules for higher dimension couplings are provided by symmetries of a compactified space as well as gauge symmetries. In refs.[5, 6], selection rules for non-renormalizable couplings are discussed within the framework of orbifold models [7]. These selection rules could lead to realistic quark mass matrices.

In this paper we discuss left-right symmetric quark mass matrices whose up- and down-sectors have the same structure. We show examples leading to these types of the realistic quark mass matrices. The μ -term is also discussed.

In general, the underlying theory like supergravity or superstring theory includes nonrenormalizable couplings like $q_{(u,d)j} Q_i H_{2,1} (\theta/M_{2,1})^{n_{ij}}$, where $q_{(u,d)j}$ denotes the SU(2) singlet quark fields and $H_{2,1}$ are the Higgs fields for the up- and down-sectors. After the field θ develops vacuum expectation value (VEV), this coupling works as the Yukawa coupling with the suppression factor $\varepsilon_{u,d}^{n_{ij}} = (\langle \theta \rangle / M_{2,1})^{n_{ij}}$. That leads to a hierarchical structure in the mass matrices and the structure depends on underlying symmetries. In general we can expect $\varepsilon_u \neq \varepsilon_d$. For example a mixing between light and

heavy Higgs fields leads to $\varepsilon_u \neq \varepsilon_d$ [2, 3].

Here we consider the following left-right symmetric mass matrices:

$$M_{u,d} = d_{u,d} \begin{pmatrix} 0 & \varepsilon_{u,d}^{n_{12}} & \varepsilon_{u,d}^{n_{13}} \\ \varepsilon_{u,d}^{n_{12}} & \varepsilon_{u,d}^{n_{22}} & \varepsilon_{u,d}^{n_{23}} \\ \varepsilon_{u,d}^{n_{13}} & \varepsilon_{u,d}^{n_{23}} & 1 \end{pmatrix}. \quad (1)$$

Note that the powers n_{ij} of the up-sector are same as those of the down-sector. The (1,2), (2,2) and (2,3) elements should be non-vanishing elements in order to lead to realistic mixing angles [8].

At first we consider the case where $\varepsilon_{u,d}^{n_{13}}$ is negligible. In this case we have two types of the mass matrices leading to the geometrical hierarchy $m_3 m_1 \approx m_2^2$ as follows,

$$3\ell = n_{12} = 3n_{23}, \quad n_{22} \geq 2n_{23}, \quad (2a)$$

$$6\ell = 2n_{12} = 3n_{22}, \quad n_{22} \leq 2n_{23}. \quad (2b)$$

We refer to these types, (2a) and (2b), as Type 1 and 2, respectively. Both types lead to the eigenvalues of masses as $m_3 = O(d_{u,d})$, $m_2 = O(\varepsilon_{u,d}^{2\ell} d_{u,d})$ and $m_1 = O(\varepsilon_{u,d}^{4\ell} d_{u,d})$. In general these types of the quarks mass matrices:

$$M_{u,d} = d_{u,d} \begin{pmatrix} 0 & a_{u,d} \varepsilon_{u,d}^{3\ell} & 0 \\ a_{u,d} \varepsilon_{u,d}^{3\ell} & b_{u,d} \varepsilon_{u,d}^m & c_{u,d} \varepsilon_{u,d}^n \\ 0 & c_{u,d} \varepsilon_{u,d}^n & 1 \end{pmatrix}, \quad (3)$$

lead to the following CKM matrix:

$$V_{\text{CKM}} \approx \begin{pmatrix} 1 - \frac{1}{2}(a_d \varepsilon_d^{3\ell} / b'_d)^2 & a_d \varepsilon_d^{3\ell} / b'_d & -a_u c_d \varepsilon_u^{3\ell} \varepsilon_d^n / b'_u \\ -a_d \varepsilon_d^{3\ell} / b'_d & 1 - \frac{1}{2}(a_d \varepsilon_d^{3\ell} / b'_d)^2 - \frac{1}{2}(c_d \varepsilon_d^n)^2 & c_d \varepsilon_d^n \\ a_d c_d \varepsilon_d^{3\ell} \varepsilon_d^n / b'_d & -c_d \varepsilon_d^n & 1 - \frac{1}{2}(c_d \varepsilon_d^n)^2 \end{pmatrix}, \quad (4)$$

where $b'_{u,d} = b_{u,d} \varepsilon_{u,d}^m - c_{u,d}^2 \varepsilon_{u,d}^{2n}$. We use the relation $\varepsilon_u < \varepsilon_d$ to get (4). Eq.(3) with $n = \ell$ corresponds to Type 1, while Type 2 is obtained by $m = 2\ell$. These types correspond to generalization of the Fritzsch ansatz [9].

Some phenomenological aspects of generic forms without the (1,1) and (1,3) elements are discussed.* The presence of the (1,3) element does not change the eigenvalues of the masses if $n_{13} \gtrsim n_{23}$ for Type 1 and $n_{13} \gtrsim n_{22}$ for Type 2. We shall not discuss the case where the (1,3) element is dominant.

The orbifold construction is one of the simplest and most interesting constructions to derive four-dimensional string vacua [7]. In orbifold models, string states consist of the bosonic string on the four-dimensional space-time and a six-dimensional orbifold, their right-moving superpartners and left-moving gauge parts. The right-moving fermionic parts are bosonized and momenta of bosonized fields span an $SO(10)$ lattice. An orbifold is obtained through a division of a six-dimensional space R^6 by a six-dimensional lattice and its automorphism θ . Closed strings on the orbifold are classified into untwisted and twisted sectors. For the θ^k -twisted sector T_k , the string coordinate satisfies the following boundary condition:

$$x_\nu(\sigma = 2\pi) = \theta^k x_\nu(\sigma = 0) + e_\nu, \quad (5)$$

where e_ν is a lattice vector. A zero-mode of this string satisfies the same condition as (5) and it is called a fixed point. The fixed point is represented in terms of the space group element (θ^k, e_ν) . All fixed points in T_k are not fixed under θ . To obtain θ -eigenstates, we have to take linear combinations of states corresponding to fixed points [11, 12]. These linear combinations have eigenvalues $\gamma = \exp[2\pi im/k]$. We take a complex basis (X_i, \overline{X}_i) ($i = 1 \sim 3$) for the compactified space, e.g. $X_1 = x_1 + ix_2$. Oscillated states in T_k are created by $\partial X_{i(k)}$ and $\partial \overline{X}_{i(k)}$.

Couplings are calculated by using vertex operators V_a corresponding to states [13, 14]. Nonvanishing couplings are invariant under a symmetry of each part of string states. Coupling terms are allowed if they are gauge invariant and space group invariant.[†] In addition the $SO(10)$ momentum

* See e.g. ref.[10].

[†] For the selection rule due to the space group, see in detail ref.[12].

should be conserved and a product of eigenvalues γ_a should satisfy $\prod_a \gamma_a = 1$. Further the corresponding correlation function $\langle V_1 \cdots V_n \rangle$ should be invariant under a Z_N rotation of oscillators as

$$\partial X_{i(k)} \rightarrow e^{2\pi i k v^i} \partial X_{i(k)}, \quad (6)$$

where $e^{2\pi i v^i}$ are eigenvalues of θ in the complex basis (X_i, \bar{X}_i) . Note that vertex operators corresponding to non-oscillated states includes the oscillators except the -1 or $-1/2$ picture [13]. These selection rules are discussed explicitly in ref.[6].

For example, the Z_6 -II orbifold has eigenvalues $v_i = (2, 1, -3)/6$. For T_1 , T_2 and T_4 , there are three fixed points on the first plane. These fixed points are denoted as $(\theta^k, i e_1)$ with $i = 0, 1, 2$. Couplings including T_4 are allowed in the following forms:

$$T_1^{2\ell} T_4^m, \quad T_3^{2\ell} T_4^{3m}, \quad T_1^{2\ell} T_2^m T_4^n, \quad T_1^\ell T_3^m T_2^p T_4^q, \quad (7)$$

where $\ell > 0, m > 0, n > 0$ and p or $q > 0$.[‡] The first coupling is allowed if $\ell + 2m = 9i + 3$ and $2\ell + m = 9j + 3$. The third and fourth couplings are forbidden unless they are point group invariant. Further the third coupling should satisfy that $\ell = 2p + 1$ and $2\ell + 2m + n = 3i$. This orbifold construction allows only the $T_1^2 T_4$ and $T_1 T_2 T_3$ couplings as renormalizable couplings of the twisted sectors.

Here we study models to lead to (2.a,b) using the above selection rules. We restrict ourselves to the simple cases where $n_{12} = 3$, i.e. $\ell = 1$ in (2.a,b) and (3). In order to obtain a left-right symmetric mass matrix, it is simple to assign one generation of the left- and right-handed quarks to the same twisted sector. We assign the Higgs fields $H_{2,1}$ to T_4 and third generations of the quark fields to T_1 so that the renormalizable coupling $T_1^2 T_4$ leads to the (3,3) element. Only the $T_1 T_3 T_4^2$ coupling induce the Yukawa coupling with

[‡]The Z_6 -II orbifold models allow other couplings. See in detail ref.[6].

the factor ε , while the couplings with ε^2 are generated by $T_1^2 T_2 T_4^2$, $T_1 T_2^2 T_3 T_4$, $T_1^2 T_3^2 T_4$ and $T_3^2 T_4^3$ couplings. In addition $T_1^2 T_2^3 T_4$, $T_1 T_2 T_3 T_4^3$, $T_1^3 T_2 T_3 T_4$ and $T_1 T_3^3 T_4^2$ couplings lead to the Yukawa couplings with ε^3 . We have to assign the second generation to T_3 or T_4 to obtain the (2,3) element of Type 1 as ε . Further we need the θ field with the VEV in T_4 and T_3 if the second generation belongs to T_3 and T_4 , respectively. In both cases, the (2,2) element is obtained as ε^2 through the $T_3^2 T_4^3$ coupling. If we assign the first generation to T_1 for the later case, the $(T_1 T_4^2) T_3^3$ coupling provides the (1,2) element as ε^3 . In this case we can assign the fixed points of the first and third generations and the Higgs fields to forbid the (1,1) and/or (1,3) elements,[§] although without this selection rule due to the space group these elements are allowed as the renormalizable coupling, $T_1^2 T_4$. Thus we obtain the following mass matrices:

$$M_{u,d} = d_{u,d} \begin{pmatrix} 0 & \varepsilon_{u,d}^3 & 0 \\ \varepsilon_{u,d}^3 & \varepsilon_{u,d}^2 & \varepsilon_{u,d} \\ 0 & \varepsilon_{u,d} & 1 \end{pmatrix}. \quad (8)$$

This corresponds to both of the types and is obtained in refs.[3, 4] by using an extra U(1) symmetry. We can obtain the similar mass matrices in the case where the first and second generations are assigned to T_1 and T_3 , respectively and θ fields in T_3 and T_4 develop VEVs. In this case it is hard to forbid completely the (1,1) and (1,3) elements. Each of these elements is induced through the $(T_1^2 T_4) T_3^2 T_4^3$ or $(T_1^2 T_4) T_4^9$ couplings, although this factor is enough suppressed. We can easily obtain the mass matrices with the suppressed (1,1) and (1,3) elements, if we assign the first and third generations to the same twisted sectors. It is hard to derive Type 1 with more suppressed (2,2) element, i.e. $m > 2$ in the Z_6 -II orbifold models, because the (2,2) element is naturally induced by the $T_3^2 T_4^3$ coupling.

Next we discuss examples leading to Type 2 with $n > 1$. It is easy to derive the (2,2) element as ε^2 for any assignment of the second generation.

[§]If the fixed points of T_1 and T_4 on the first plane do not satisfy the space group invariance, one cannot generate these elements by the θ field with the VEV in T_3 .

However, it is hard to obtain the (1,2) element as ε^3 when the second generation is assigned to T_1 . If the second generation belongs to T_4 , the matrix (8) is naturally obtained as above. Thus we consider the case where the second generation is assigned to T_3 . Further we assign the first generation to T_2 . Suppose that θ fields in T_1 , T_2 and T_4 develop VEVs. Then we obtain the (2,2) element as ε^2 through the $(T_3^2 T_4) T_4^2$ coupling. In this case the (1,1), (1,2) and (2,3) element are provided as ε^3 , ε^2 and ε by the $(T_2^2 T_4)(T_1^2 T_2)$, $(T_2 T_3 T_4)(T_1 T_2)$ and $(T_1 T_3 T_4) T_4$ coupling without taking into account other selection rules. We assume that the θ fields with the VEVs in T_2 and T_4 have $\gamma = -1$, while the other fields have $\gamma = 1$. Then the above couplings are forbidden. Instead the $(T_2^2 T_4)(T_1^6 T_4^2)$, $(T_2 T_3 T_4) T_1^3$ and $(T_1 T_3 T_4) T_2^2$ couplings generate the (1,1), (1,2) and (2,3) elements as ε^8 , ε^3 and ε^2 , respectively. Then we obtain the following quark mass matrix:

$$M_{u,d} = d_{u,d} \begin{pmatrix} \varepsilon_{u,d}^8 & \varepsilon_{u,d}^3 & \varepsilon_{u,d}^3 \\ \varepsilon_{u,d}^3 & \varepsilon_{u,d}^2 & \varepsilon_{u,d}^2 \\ \varepsilon_{u,d}^3 & \varepsilon_{u,d}^2 & 1 \end{pmatrix}, \quad (9)$$

where the (1,3) element is obtained through the $(T_1 T_2 T_4)(T_1 T_2^2)$ coupling. We can obtain more suppressed (1,3) element if we take into account the selection rule due to the space group invariance. For example we assign the fixed point of the T_1 field with the VEV to (θ, e_1) on the first plane, while the fixed point of the other fields is assigned to the origin $(\theta^k, 0)$. Then the (1,3) element is derived as ε^9 through the $(T_1 T_2 T_4) T_1^9$ coupling, while the $(T_2 T_3 T_4) T_1^3$ coupling of the (1,2) element is allowed. Other assignment of the fixed points can lead to similar mass matrices. We neglect the (1,1) and (1,3) elements to obtain the following mass matrices:

$$M_{u,d} = d_{u,d} \begin{pmatrix} 0 & \varepsilon_{u,d}^3 & 0 \\ \varepsilon_{u,d}^3 & \varepsilon_{u,d}^2 & \varepsilon_{u,d}^2 \\ 0 & \varepsilon_{u,d}^2 & 1 \end{pmatrix}. \quad (10)$$

Eq.(10) is one of the simplest mass matrices [15]. Because four texture zeros are included totally for the up- and down-sectors and for each matrix there

appear only three hierarchies, 1, ε^2 and ε^3 . This type can lead to the realistic mixing angles (4). In this case the mixing angles satisfy the following relations:

$$|V_{us}|^2 \approx |V_{cb}| \approx \frac{m_d}{m_s}, \quad \frac{|V_{ub}|}{|V_{cb}|} \approx \sqrt{\frac{m_u}{m_c}}, \quad \frac{|V_{td}|}{|V_{ts}|} \approx \sqrt{\frac{m_d}{m_s}}. \quad (11)$$

We can derive realistic mass matrices similar to (9) and (10) in the case where the second generation is assigned to T_2 . In this case we need several fields with VEVs.

We can derive Type 2 with $n > 2$. For example we assign the first and second generations to T_1 and T_2 , respectively. Suppose that θ fields in T_1 and T_3 develop VEVs. Then we obtain the same mass matrices as (10) except the (2,3) element replacing ε^3 . Further we can derive Type 2 with $n > 3$, but the larger values of n do not lead to the realistic mixing angles (4).

We have derived some examples of Type 1 and 2 (2a,b) with $n_{12} = 3$. Similarly we could obtain these types of the quark mass matrices with $n_{12} = 3\ell$ ($\ell > 1$), although in some cases it is hard to rule out lower dimension operators. It is also interesting to derive the quark mass matrices where the geometrical hierarchy is obtained approximately, e.g. the case with $3n_{12} = 4n_{22}$ and $n_{23} \geq n_{22}$. This example corresponds to some down-sectors in ref.[2].

World-sheet instantons lead to another suppression factor as $\exp[-a_{ij}T]$, where T is the moduli parameter and a_{ij} is a constant [14]. These effects provide the coefficients like (3). Also CP phases are induced to some elements in the case with nonvanishing background anti-symmetric tensors [16], although in some orbifold models the physical CP phase do not appear in the limit $\varepsilon_{u,d} \rightarrow 0$ [6].[¶] The suppression factor $\exp[-a_{ij}T]$ and the CP phase of each element are in general different from those of other elements. The difference between d_u and d_d can be explained by the difference of the VEVs of $H_{2,1}$

[¶] See also ref.[17].

or the world-sheet instanton effects.^{||} Further, we have to estimate radiative corrections from a high energy to a low energy to compare the prediction with experiment. However, the hierarchical structure, i.e. the powers n_{ij} are not changed [18, 2], while the coefficients in (3) are changed by $O(1)$.

At last we discuss the μ -term. In the above examples, both of the Higgs fields, $H_{2,1}$ are assigned to T_4 . Thus the μ -term is forbidden as a renormalizable couplings. After the symmetry breaking, some high dimension operators could generate the effective μ -term, which is naturally suppressed. For instance, the μ -term in the example leading to (10) should include $T_1^6 T_2^2 T_4^2$ coupling.

To sum up, We have studied on left-right symmetric quark mass matrices, which have the same structure for the up- and down-sectors. We have shown examples leading to these type of the mass matrices within the framework of the orbifold models. For the Z_6 -II orbifold models, it is hard to obtain Type 1 except (8). Eq.(10) is one of the simplest and most realistic quark mass matrices, which cannot be derived by assuming an extra U(1) symmetry. The above analyses constrain how to assign the quark fields to the twisted sectors. That is very useful to model building.

Acknowledgement

The author would like to thank Z. Xing for numerous useful discussions and he also thanks C.S. Lim, H. Nakano and D. Suematsu.

References

- [1] C.D. Froggatt and H.B. Nielsen, Nucl. Phys. **B147** (1979) 277.

^{||}We could also explain this difference by assuming an extra U(1) symmetry.

- S. Dimopoulos, Phys. Lett. **B129** (1983) 417.
- M. Leurer, Y. Nir and N. Seiberg, Nucl. Phys. **B398** (1993) 319;
Nucl. Phys. **B420** (1994) 468.
- Y. Nir and N. Seiberg, Phys. Lett. **B309** (1993) 337.
- E. Papageorgiu, Z. Phys. **C64** (1994) 509.
- E. Dudas, S. Pokorski and C.A. Savoy, preprint SPhT Saclay T95/027
(hep-ph/9504292).
- [2] P. Ramond, R.G. Roberts and G.G. Ross, Nucl. Phys. **B406** (1993) 19.
- [3] L.E. Ibáñez and G.G. Ross, Phys. Lett. **B332** (1994) 100.
- [4] P. Binétruy and P. Ramond, Phys. Lett. **B350** (1995) 49.
- [5] M. Cvetič, Phys. Rev. Lett. **59** (1987) 1795; Phys. Rev. Lett. **59** (1987) 2829.
- A. Font, L.E. Ibáñez, H.P. Nilles and F. Quevedo, Nucl. Phys. **B307** (1988) 109; Phys. Lett. **B210** (1988) 101; Phys. Lett. **B213** (1988) 274.
- A. Font, L.E. Ibáñez, F. Quevedo and A. Sierra, Nucl. Phys. **B331** (1990) 421.
- [6] T. Kobayashi, preprint LMU-TPW 95-5 (hep-ph/9504371) to be published in Phys. Lett. **B**.
- [7] L. Dixon, J. Harvey, C. Vafa and E. Witten, Nucl. Phys. **B261** (1985) 678; Nucl. Phys. **B274** (1986) 285.
- L.E. Ibáñez, J. Mas, H.P. Nilles and F. Quevedo, Nucl. Phys. **B301** (1988) 157.
- Y. Katsuki, Y. Kawamura, T. Kobayashi, N. Ohtsubo, Y. Ono and K. Tanioka, Nucl. Phys. **B341** (1990) 611.
- [8] L.J. Hall and A. Rasin, Phys. Lett. **B315** (1993) 164.

- [9] H. Fritzsch, Phys. Lett. **B73** (1978) 317; Nucl. Phys. **B155** (1979) 189.
- [10] H. Fritzsch and Z. Xing, preprint MPI-PhT/95-08 (hep-ph/9502297) to be published in Phys. Lett. **B**.
- [11] T. Kobayashi and N. Ohtsubo, Phys. Lett. **B245** (1990) 441.
- [12] T. Kobayashi and N. Ohtsubo, Int. J. Mod. Phys. **A9** (1994) 87.
- [13] D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986) 93.
- [14] S. Hamidi and C. Vafa, Nucl. Phys. **B279** (1987) 465.
L. Dixon, D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B282** (1987) 13.
- [15] Z. Xing, private communication.
- [16] J. Erler, D. Jungnickel and J. Lauer, Phys. Rev. **D45** (1992) 3651.
D. Jungnickel, J. Lauer, M. Spaliński and S. Stieberger, Mod. Phys. Lett. **A7** (1992) 3059.
J. Erler, D. Jungnickel, M. Spaliński and S. Stieberger, Nucl. Phys. **B397** (1993) 379.
- [17] C.S. Lim, Phys. Lett. **B256** (1991) 233.
M. Dine, R.G. Leigh and D.A. MacIntire, Phys. Rev. Lett. **69** (1992) 2030.
K. Choi, D.B. Kaplan and A.E. Nelson, Nucl. Phys. **B391** (1993) 515.
T. Kobayashi and C.S. Lim, Phys. Lett. **B343** (1995) 122.
- [18] M. Olechowski and S. Pokorski, Phys. Lett. **B257** (1991) 388.